



# Cardiomyopathy

ICD-10-CM

Clinical overview

## Definition

Cardiomyopathy is a disease of the heart muscle that impairs the function of the heart.

## Types

Cardiomyopathy can be classified as primary or secondary and ischemic or nonischemic.

- **Primary cardiomyopathy** is a noninflammatory disease of the heart muscle, often of obscure or unknown cause, that occurs in the absence of other cardiac conditions or systemic disease processes.
- **Secondary cardiomyopathy** is caused by a known medical condition (such as hypertension, valve disease, congenital heart disease or coronary artery disease).
- **Ischemic cardiomyopathy** is caused by coronary artery disease and heart attacks, which result in lack of blood flow to the heart muscle, thereby causing damage to the heart muscle.
- **Nonischemic cardiomyopathy** is a type of cardiomyopathy not related to coronary artery disease or poor coronary artery blood flow. There are three main types of nonischemic cardiomyopathy:
  - **Dilated cardiomyopathy (also known as congestive cardiomyopathy)** – This is the most common type of cardiomyopathy. In this disorder, the heart's main pumping chamber – the left ventricle – becomes enlarged (dilated), its pumping ability becomes less forceful and blood doesn't flow as easily through the heart.
  - **Hypertrophic cardiomyopathy** – This type involves abnormal growth or thickening of the heart muscle, particularly affecting the muscle of the left ventricle. As thickening occurs, the heart tends to stiffen and the size of the pumping chamber may shrink, interfering with the heart's ability to deliver blood to the body.
  - **Restrictive cardiomyopathy** – The heart muscle in people with restrictive cardiomyopathy becomes rigid and less elastic, meaning the heart can't properly expand and fill with blood between heartbeats.

Some cardiomyopathies can be reversible. For example:

- Alcoholic cardiomyopathy sometimes can be reversed with complete cessation of alcohol intake.
- Takotsubo cardiomyopathy is a reversible, stress-induced cardiomyopathy.

## Causes

The cause is usually unknown (primary cardiomyopathy), although contributing factors sometimes can be identified. Some of the possible known causes include:

- Long-term high blood pressure
- Coronary artery disease
- Heart valve problems
- Chronic rapid heart rate
- Certain viral infections
- Some chemotherapy drugs
- Pregnancy
- Excessive, long-term use of alcohol
- Heart damage, due to a previous heart attack
- Metabolic disorders (thyroid disease, diabetes, etc.)
- Nutritional deficiencies of essential vitamins and minerals
- Abuse of cocaine or antidepressant medications
- Hemochromatosis – disorder in which iron is not properly metabolized, causing build-up in various organs, including heart muscle. This can lead to a weakening of the heart muscle, resulting in dilated cardiomyopathy.

## Signs and symptoms

There may be no signs or symptoms in the early stages of the disease. But as the condition advances, signs and symptoms usually appear and may include:

- Shortness of breath, especially with physical exertion
- Swelling of lower extremities, abdomen and neck veins
- Fatigue
- Chest pain
- Irregular heartbeats
- Heart murmurs
- Dizziness and lightheadedness
- Fainting

## Possible complications

- Heart failure
- Blood clots
- Heart valve problems with associated murmurs
- Cardiac arrest and sudden death



# Cardiomyopathy

ICD-10-CM

Clinical overview

## Diagnostic tools

- Medical history and physical exam
- Blood tests, such as measurement of brain natriuretic peptide (BNP) – a protein produced in the heart for which the blood level increases when the heart is under stress from heart failure, a common complication of cardiomyopathy
- Chest X-ray (to check for signs such as enlarged heart or fluid buildup in lungs)
- Electrocardiogram (ECG or EKG)
- Holter monitoring and cardiac event recording
- Echocardiogram
- Cardiac stress testing
- Cardiac magnetic resonance imaging (MRI)
- Cardiac catheterization and heart biopsy

## Treatment

- Lifestyle changes:
  - Heart-healthy diet
  - Weight control
  - Stress management
  - Physical activity and exercise
  - Smoking cessation
- Medications:
  - Blood thinners to prevent clots
  - Antiarrhythmics to control heart rate and rhythm
  - Antihypertensives for blood pressure control
  - Diuretics (“water pills”) to remove excess sodium and reduce excess fluid in the blood
- Nonsurgical procedure:
  - Alcohol septal ablation, in which a type of alcohol (ethanol) is injected through a tube into the small artery that supplies blood to the thickened area of heart muscle. The alcohol shrinks the thickened heart tissue to a more normal size, allowing blood to flow freely through the ventricle of the heart, which results in improved symptoms.
- Cardiac device implantation:
  - Pacemaker
  - Cardioverter-defibrillator
  - Left ventricular assist device
- Surgical procedures:
  - Heart transplant (a last resort for severe, end-stage cardiomyopathy that cannot be controlled by other means)



# Cardiomyopathy

ICD-10-CM

Best documentation practices for physicians

## Subjective

The subjective section of the office note should document current related patient complaints and symptoms. If there are none, the office note should show the patient was screened for current related complaints or symptoms.

## Objective

In the objective section, include associated physical exam findings (such as edema/swelling of the lower extremities, abdomen or neck veins) and related diagnostic testing results.

## Assessment

### Specificity:

- The term “cardiomyopathy” is broad and nonspecific. It is important to describe the particular type of cardiomyopathy to the highest level of specificity.
- Document the current status of cardiomyopathy (stable, improved, worsening, etc.).

### Abbreviations:

- A good rule of thumb for any medical record is to limit – or avoid altogether – the use of abbreviations. There are several commonly used medical abbreviations for different types of cardiomyopathy (CM, CMP, HCM, HOCM, etc.), but some of these abbreviations have other meanings. The meaning of an abbreviation can often be determined based on context, but this is not always true.
- Best practice is to always document the specific type of cardiomyopathy by spelling it out in full.

### Associated conditions:

Clearly link secondary cardiomyopathy to the underlying causative condition by using terms such as “due to,” “secondary to,” “associated with,” “related to,” etc.

### Current versus historical/transient:

- Do not use the descriptor “history of” to describe current cardiomyopathy. In diagnosis coding, “history of” means the condition occurred in the past and is no longer a current problem.
- Temporary or transient cardiomyopathy that occurred in the past and is no longer present should not be documented as if it is current.

## Plan

- Document a specific and concise treatment plan.
- Clearly link the cardiomyopathy diagnosis to any medications being used to treat the condition.
- Document referrals to specialists or other providers.
- Include the date of the patient’s next appointment.

## Electronic health record (EHR) issues

### Other and unspecified codes with descriptions:

Some electronic health records (EHRs) insert ICD-10-CM codes with corresponding descriptions into the assessment section of the office note rather than a provider-stated final diagnosis. For example:

I42.8 Other cardiomyopathies

I42.9 Cardiomyopathy, unspecified

These are vague descriptions and incomplete diagnoses.

- Codes titled “other” or “other specified” are for use when the medical record provides a specific diagnosis description for which a specific code does not exist.
- The “other” ICD-10-CM code with description should not be used, by itself, as a final diagnosis without clear documentation that specifies the particular “other” condition.
- Unspecified diagnosis descriptions should be used only when sufficient clinical information is not known or available to the provider at the time of the encounter.

### Mismatch between final diagnostic statement and EHR-inserted diagnosis code with description:

Another scenario that causes confusion is one in which the assessment section documents a provider-stated diagnosis *PLUS* an EHR-inserted diagnosis code with description that does not match – or may even contradict – the stated diagnosis. Example:

#### Assessment: Ischemic cardiomyopathy

I42.Ø Dilated cardiomyopathy

Here the final diagnosis in **bold** in the Assessment is “Ischemic cardiomyopathy”, which codes to I25.5.

The EHR-inserted diagnosis code with description that follows, however, is I42.Ø, Dilated cardiomyopathy.



# Cardiomyopathy

ICD-10-CM

Best documentation practices for physicians

To avoid confusion and ensure accurate diagnosis code assignment, the provider-stated final diagnosis must either

- a) match the code with description; OR
- b) it must classify in ICD-10-CM to the EHR-inserted diagnosis code with description.

**Note:** ICD-10-CM is a statistical classification; it is not a substitute for a healthcare provider's final diagnostic statement. It is the healthcare provider's responsibility to provide legible, clear, concise and specific documentation of each final diagnosis described to the highest level of specificity, which is then translated to a code for reporting purposes. It is not appropriate for healthcare providers to simply list a code number or select a code number from a list of codes in place of a written final diagnosis.

Reference: AHA Coding Clinic, Code Number in Lieu of a Diagnosis, Fourth quarter 2015, pages 34-35



# Cardiomyopathy

ICD-10-CM

Tips and resources for coders

## Coding basics

To ensure accurate and specific diagnosis code assignment, the coder must review the entire medical record and then, in accordance with ICD-10-CM official coding conventions and guidelines:

- Search the alphabetic index for the specific diagnosis description and the corresponding code.
- Verify the code in the tabular list, carefully following all instructional notes.

## Coding cardiomyopathy

Many of the most common cardiomyopathies classify to category **I42**, Cardiomyopathy.

- A fourth character is required to specify the particular type of cardiomyopathy.
- The broad and nonspecific final diagnosis of “cardiomyopathy” leads to the broad and nonspecific diagnosis code **I42.9**, Cardiomyopathy, unspecified.
- Code **I42.9** should be assigned only when no information in the medical record identifies the particular type of cardiomyopathy.

Hypertensive cardiomyopathy classifies to category **I11**, Hypertensive heart disease, with an additional code of **I43**, Cardiomyopathy in diseases classified elsewhere.

Congestive cardiomyopathy is also known as dilated cardiomyopathy. Both of these descriptions classify to code **I42.0**, Dilated cardiomyopathy.

- Congestive cardiomyopathy often is associated with congestive heart failure and basically has the same symptoms.
- Treatment typically focuses on management of the congestive heart failure; therefore, heart failure (category **I50**.-) is reported as the principal diagnosis with an additional code for the cardiomyopathy.

Hypertrophic cardiomyopathy can be obstructive or nonobstructive.

- **I42.1** Obstructive hypertrophic cardiomyopathy
- **I42.2** Other hypertrophic cardiomyopathy  
Includes nonobstructive hypertrophic cardiomyopathy

## Reminders

- Use caution when coding cardiomyopathy from abbreviations (CM, HCM, HOCM, etc.). A code should not be assigned unless the meaning of the abbreviation is clear based on review of the entire medical record.
- Watch for modifying descriptors that affect code assignment (secondary, alcoholic, nutritional, metabolic or cardiomyopathy due to other diseases).
- Some secondary cardiomyopathies are coded with a single combination code, while other secondary cardiomyopathies require the use of two codes. See coding examples on page 6.
- Takotsubo cardiomyopathy is a reversible form of cardiomyopathy that classifies to code **I51.81**, Takotsubo syndrome. This code includes the following conditions:
  - Reversible left ventricular dysfunction following sudden emotional stress
  - Stress-induced cardiomyopathy
  - Takotsubo cardiomyopathy
  - Transient left ventricular apical ballooning syndrome
- The term “ischemic cardiomyopathy” is sometimes used to refer to a condition in which ischemic heart disease causes diffuse fibrosis or multiple infarction, leading to heart failure with left ventricular dilation. This is not a true cardiomyopathy. When no further clarification is available, this condition is coded to **I25.5**, Ischemic cardiomyopathy.
- In ICD-10-CM, dilated cardiomyopathy and ischemic cardiomyopathy are classified to different codes. When a medical record documents a current diagnosis of ischemic dilated cardiomyopathy, both codes are needed to fully capture this condition:
  - **I25.5** Ischemic cardiomyopathy
  - **I42.0** Dilated cardiomyopathy
- Nonischemic cardiomyopathy with no other description and no mention of cause codes to **I42.8**.

## ALPHABETIC INDEX

**Cardiomyopathy** (familial) (idiopathic)  
nonischemic – *see also* by cause **I42.8**

## TABULAR LIST

**I42.8** Other cardiomyopathies



# Cardiomyopathy

ICD-10-CM

Tips and resources for coders

## Coding examples

Example 1	
Final diagnosis	Cardiomyopathy (nonspecific)
ICD-10-CM code(s)	I42.9 Cardiomyopathy, unspecified

  

Example 2	
Final diagnosis	Ischemic dilated cardiomyopathy
ICD-10-CM code(s)	I25.5 Ischemic cardiomyopathy I42.0 Dilated cardiomyopathy

  

Example 3	
Final diagnosis	Hypertensive cardiomyopathy
ICD-10-CM code(s)	I11.9 Hypertensive heart disease without heart failure I43 Cardiomyopathy in disease classified elsewhere

  

Example 4	
Final diagnosis	Cardiomyopathy due to sarcoidosis
ICD-10-CM code(s)	D86.85 Sarcoid myocarditis

  

Example 5	
Final diagnosis	Metabolic cardiomyopathy
ICD-10-CM code(s)	E88.9 Metabolic disorder, unspecified I43 Cardiomyopathy in diseases classified elsewhere

  

Example 6	
Final diagnosis	Rheumatic cardiomyopathy
ICD-10-CM code(s)	I09.0 Rheumatic myocarditis

  

Example 7	
Final diagnosis	Stress-induced cardiomyopathy
ICD-10-CM code(s)	I51.81 Takotsubo syndrome

  

Example 8	
Final diagnosis	Restrictive cardiomyopathy
ICD-10-CM code(s)	I42.5 Other restrictive cardiomyopathy

  

Example 9	
Final diagnosis	End-stage dilated cardiomyopathy
ICD-10-CM code(s)	I42.0 Dilated cardiomyopathy

  

Example 10	
Final diagnosis	Alcoholism in remission with alcoholic cardiomyopathy
ICD-10-CM code(s)	F10.21 Alcohol dependence, in remission I42.6 Alcoholic cardiomyopathy

**References:** American Hospital Association Coding Clinic; ICD-10-CM Official Guidelines for Coding and Reporting; ICD-10-CM and ICD-10-PCS Coding Handbook; Mayo Clinic; National Heart, Lung, and Blood Institute