

Remove above portion before faxing. Please complete the prescription form in its entirety and fax with secure cover sheet to the number above.

Hemophilia and Bleeding Disorders Prescription Form

Patient information

Patient: _____ Female Male DOB: _____ Insurance plan: _____ Plan ID #: _____
 Address: _____ City: _____ State: _____ ZIP code: _____
 Home phone #: _____ Cell phone #: _____ Caregiver: _____ Caregiver phone #: _____
 Other medical conditions: _____ Allergies: No Yes: _____
 Height: _____ Weight: _____ lb kg Date: _____ ICD-10 code: _____ Diagnosis: _____ Date: _____

Clinical information

<input type="checkbox"/> Hereditary Factor VIII Deficiency: D66	Severity: <input type="checkbox"/> Mild <input type="checkbox"/> Moderate <input type="checkbox"/> Severe	Planned procedure: <input type="checkbox"/> Dental extraction(s) Date(s): _____ <input type="checkbox"/> Surgical procedure(s) Date(s): _____
<input type="checkbox"/> Hereditary Factor IX Deficiency: D67	Target joint(s)? <input type="checkbox"/> No <input type="checkbox"/> Yes: _____	
<input type="checkbox"/> Hereditary Factor XI Deficiency: D68.1	Venous access: <input type="checkbox"/> PIV <input type="checkbox"/> PICC <input type="checkbox"/> Port	
<input type="checkbox"/> Von Willebrand's disease: D68.0	<input type="checkbox"/> Central line, type: _____	
<input type="checkbox"/> Acquired hemophilia: D68.311	Inhibitor? <input type="checkbox"/> No <input type="checkbox"/> Yes: _____ BU	
<input type="checkbox"/> _____	Doses to keep on hand: _____	

Prescription information

Note: Ohio law allows one prescription per preprinted order form. Please use additional forms for more than one prescription.

<input type="checkbox"/> Advate	<input type="checkbox"/> BeneFix	<input type="checkbox"/> Helixate FS	<input type="checkbox"/> Jivi	<input type="checkbox"/> Novoeight	<input type="checkbox"/> Recombinate
<input type="checkbox"/> Adynovate	<input type="checkbox"/> Corifact	<input type="checkbox"/> Hemlibra	<input type="checkbox"/> Kcentra	<input type="checkbox"/> Nuwiq	<input type="checkbox"/> Tretten
<input type="checkbox"/> Afstyla	<input type="checkbox"/> Elocrate	<input type="checkbox"/> Hemofil M	<input type="checkbox"/> Koate-DVI	<input type="checkbox"/> Profilnine SD	<input type="checkbox"/> Vonvendin
<input type="checkbox"/> Alphanate	<input type="checkbox"/> Esperoct	<input type="checkbox"/> Humate-P	<input type="checkbox"/> Kovaltry	<input type="checkbox"/> Riastap	<input type="checkbox"/> Wilate
<input type="checkbox"/> AlphaNine SD	<input type="checkbox"/> Feiba FS	<input type="checkbox"/> Idelvion	<input type="checkbox"/> Lysteda	<input type="checkbox"/> Rebinyn	<input type="checkbox"/> Xyntha
<input type="checkbox"/> Alprolix	<input type="checkbox"/> Fibryga	<input type="checkbox"/> Ixinity	<input type="checkbox"/> NovoSeven RT	<input type="checkbox"/> Rixubis	
<input type="checkbox"/> Other _____					

Prophylaxis (+/- 10% or other: _____) Dose: _____ IU IU/kg RCOF Frequency: _____ # of doses: _____ Refills: _____
 Immune tolerance (+/- 10% or other: _____) Target dose: _____ IU IU/kg RCOF Frequency: _____ # of doses: _____ Refills: _____
 Breakthrough bleed (+/- 10% or other: _____) Minor dose: _____ IU IU/kg RCOF Frequency: _____ # of doses: _____ Refills: _____
 Moderate dose: _____ IU IU/kg RCOF Frequency: _____ # of doses: _____ Refills: _____
 Major dose: _____ IU IU/kg RCOF Frequency: _____ # of doses: _____ Refills: _____
 Hemlibra Initial dose: 3mg/kg _____ mg/kg mg Subsequent dose: 1.5mg/kg 3mg/kg 6mg/kg _____ mg/kg mg
 Directions: _____ Directions: _____
 # of doses: _____ Refills: _____ # of doses: _____ Refills: _____

Skilled nursing visit to establish venous access, provide patient education related to therapy and disease state, administer medication as prescribed, and assess general status and response to therapy. Frequency based on prescription dose orders.

Additional medications/supplies	Directions	Quantity	Refills
<input type="checkbox"/> Amicar <input type="checkbox"/> 500 mg tab. <input type="checkbox"/> 1000 mg tab. <input type="checkbox"/> 250 mg/mL sol.	_____	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> desmopressin 4 mcg/mL inj.	_____	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> heparin 100U/mL 5 mL PFS For central line patients.	Flush line with 5 mL, after final saline flush.	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> lidocaine 2.5% and prilocaine 2.5% cream	Apply topically to needle insertion site, 30–60 minutes prior to insertion.	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> sodium chloride 0.9% 10 mL flush	Flush line with 10 mL, before and after inf. and p.r.n. line care.	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> Stimate 1.5 mg/mL sol.	_____	<input type="checkbox"/> _____	<input type="checkbox"/> _____
<input type="checkbox"/> tranexamic acid 650 mg tab.	_____	<input type="checkbox"/> _____	<input type="checkbox"/> _____

Prescriber and shipping information (please print)

Prescriber: _____ NPI: _____
 Ship to: Patient Office Other: _____
 Office address: _____ City: _____ State: _____ ZIP code: _____
 Office phone number: _____ Office fax number: _____
 Signature: _____ Date: _____

We will dispense this prescription as generic, unless the prescriber indicates "Dispense as Written" here: _____
 The prescriber is to comply with his/her state-specific prescription requirements, such as e-prescribing, state-specific prescription form and fax language.
 Noncompliance with state-specific requirements could result in outreach to the prescriber.