

Remove above portion before faxing. Please complete the prescription form in its entirety and fax with secure cover sheet to the number above.

**IVIG and General Immune Disorders Prescription Form**

**Patient information**

Patient: \_\_\_\_\_  Female  Male DOB: \_\_\_\_\_ Insurance plan: \_\_\_\_\_ Plan ID #: \_\_\_\_\_  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ ZIP code: \_\_\_\_\_  
 Home phone #: \_\_\_\_\_ Cell phone #: \_\_\_\_\_ Caregiver: \_\_\_\_\_ Caregiver phone #: \_\_\_\_\_  
 Other medical conditions: \_\_\_\_\_ Allergies:  No  Yes: \_\_\_\_\_  
 Height: \_\_\_\_\_ Weight: \_\_\_\_\_  lb  kg Date: \_\_\_\_\_ ICD-10 code: \_\_\_\_\_ Diagnosis date: \_\_\_\_\_

**Clinical information**

ICD-10 immunology: <input type="checkbox"/> D80.0 Congenital hypogam <input type="checkbox"/> D83.9 CVID (unspecif) <input type="checkbox"/> D81.9 SCID (unspecif) ICD-10 neurology: <input type="checkbox"/> G61.81 CIDP <input type="checkbox"/> G61.82 MMN <input type="checkbox"/> G35 MS (rel-remit) <input type="checkbox"/> G61.0 GBS <input type="checkbox"/> G70.01 MG ICD-10 rheumatology: <input type="checkbox"/> M33.20 Polymyositis <input type="checkbox"/> M33.90 Dermatopolymyositis <input type="checkbox"/> _____ Concurrent therapies: _____ Date: _____ Adverse reactions with previous IG treatments? _____ If so, what brand of IVIG caused the reaction? _____	Site of care: <input type="checkbox"/> Home <input type="checkbox"/> MDO <input type="checkbox"/> Clinic: _____ Venous access (for applicable therapies): <input type="checkbox"/> PIV <input type="checkbox"/> PICC <input type="checkbox"/> Port <input type="checkbox"/> Central line, type: _____ Infusion method (for applicable therapies): <input type="checkbox"/> Gravity <input type="checkbox"/> Pump First dose <input type="checkbox"/> No <input type="checkbox"/> Yes Expected date of first/next infusion: _____
--	---

**Prescription information** Note: Ohio law allows one prescription per preprinted order form. Please use additional forms for more than one prescription. Pharmacy to dispense ancillary supplies, as needed, to establish IV and administer drug.

Intramuscular	Subcutaneous	Intravenous	
<input type="checkbox"/> GamaSTAN S/D <input type="checkbox"/> WinRho	<input type="checkbox"/> Gammagard 10% Liquid <input type="checkbox"/> Hizentra 20% <input type="checkbox"/> Gammaked 10% <input type="checkbox"/> Vials <input type="checkbox"/> PFS <input type="checkbox"/> Gamunex-C 10% <input type="checkbox"/> Panzyga <input type="checkbox"/> Hyqvia 10% <input type="checkbox"/> Xembify	<input type="checkbox"/> Gammagard 10% Liquid <input type="checkbox"/> Gammagard S/D 5% <input type="checkbox"/> Gammagard S/D 10% <input type="checkbox"/> Gammaked 10% <input type="checkbox"/> Gamunex-C 10%	<input type="checkbox"/> Flebogamma 5% <input type="checkbox"/> Flebogamma 10% <input type="checkbox"/> Octagam 10% <input type="checkbox"/> Octagam 5% <input type="checkbox"/> Privigen 10%
Dose	Directions	Quantity	Refills
_____	_____	_____	_____

Skilled nursing visit to establish venous access, provide patient education related to therapy and disease state, administer medication as prescribed, and assess general status and response to therapy. Frequency based on prescription dose orders.

Pretreatment:	<input type="checkbox"/> acetaminophen <input type="checkbox"/> 325 mg tab. <input type="checkbox"/> 500 mg tab. <input type="checkbox"/> diphenhydramine <input type="checkbox"/> 25 mg cap. <input type="checkbox"/> 50 mg tab. <input type="checkbox"/> _____	<input type="checkbox"/> 10 cap. <input type="checkbox"/> 10 tab. <input type="checkbox"/> _____	<input type="checkbox"/> 11 <input type="checkbox"/> _____
Anaphylactic treatment:	<input type="checkbox"/> diphenhydramine <input type="checkbox"/> 50 mg/mL vial <input type="checkbox"/> 25 mg cap. <input type="checkbox"/> 50 mg tab. <input type="checkbox"/> Infuse slowly IV p.r.n. anaphylaxis. <input type="checkbox"/> Take 25–50 mg p.o. p.r.n. anaphylaxis. <input type="checkbox"/> epinephrine <input type="checkbox"/> 0.3 mg auto-injector <input type="checkbox"/> 0.15 mg auto-injector <input type="checkbox"/> Inject IM p.r.n. anaphylaxis.	<input type="checkbox"/> 1 vial <input type="checkbox"/> 10 cap. <input type="checkbox"/> 10 tab. <input type="checkbox"/> _____	<input type="checkbox"/> 0 <input type="checkbox"/> _____
<input type="checkbox"/> lidocaine 2.5% and prilocaine 2.5% cream	Apply topically to needle insertion site, 30–60 minutes prior to insertion.	<input type="checkbox"/> 28-day supply <input type="checkbox"/> _____	<input type="checkbox"/> 11 <input type="checkbox"/> _____
<input type="checkbox"/> sodium chloride 0.9% 10 mL flush	Flush line with 10 mL, before and after inf. and p.r.n. line care.	<input type="checkbox"/> 28-day supply <input type="checkbox"/> _____	<input type="checkbox"/> 11 <input type="checkbox"/> _____
<input type="checkbox"/> heparin 100U/mL 5 mL PFS - For central line patients	Flush line with 5 mL, after final saline flush.	<input type="checkbox"/> 28-day supply <input type="checkbox"/> _____	<input type="checkbox"/> 11 <input type="checkbox"/> _____
<input type="checkbox"/> Hydration fluid	<input type="checkbox"/> 0.9% NS _____ mL infused over _____ minutes Timing: <input type="checkbox"/> : _____ minutes pre-IG infusion <input type="checkbox"/> post-IG infusion <input type="checkbox"/> during IVIG infusion <input type="checkbox"/> D5W _____ mL infused over _____ minutes Timing: <input type="checkbox"/> : _____ minutes pre-IG infusion <input type="checkbox"/> post-IG infusion <input type="checkbox"/> during IVIG infusion	<input type="checkbox"/> 28-day supply <input type="checkbox"/> _____	<input type="checkbox"/> 11 <input type="checkbox"/> _____

**Prescriber and shipping information (please print)**

Prescriber: \_\_\_\_\_ NPI: \_\_\_\_\_ Office phone number: \_\_\_\_\_  
 Ship to:  Patient  Office  Other: \_\_\_\_\_ Office fax number: \_\_\_\_\_  
 Office address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ ZIP code: \_\_\_\_\_  
 Signature: \_\_\_\_\_ Date: \_\_\_\_\_  
 We will dispense this prescription as generic, unless the prescriber indicates "Dispense as Written" here: \_\_\_\_\_  
 The prescriber is to comply with his/her state-specific prescription requirements, such as e-prescribing, state-specific prescription form and fax language. Noncompliance with state-specific requirements could result in outreach to the prescriber.